

## ACAAI COVID-19 Vaccine Comparison Chart - 1/13/2021

Vaccine	Active Ingredient	Dose	Schedule	Age approved	Efficacy against COVID 19	Adverse reactions	Other Ingredients	Contra Indications
Pfizer-BioNTech	m RNA	0.3 ml	2 doses , 21 days apart	16 and up	18-64 years 95.1% 65 and above 94.7 %	Lasted 2-3 days, more adverse reactions after second dose Pain at site, Fatigue, Headache, Myalgia, Chills, Arthralgia Anaphylaxis* ( post EUA)	1.Lipids - Including Polyethylene glycol 2. Salts 3. Sugars – Sucrose No preservatives	History of allergic reactions to vaccines / vaccine ingredients including PEG , Polysorbate
Moderna	m RNA	0.5ml	2 doses , 28 days apart	18 and up	18 to < 65 years – 95.6 % >65 years– 86.4 %	Lasted 2-3 days, more adverse reactions after second dose Pain at site, Fatigue, Headache, Myalgia, Chills, Arthralgia	1.Lipids - Including Polyethylene glycol 2. Salts 3. Sugars – Sucrose No preservatives	History of allergic reactions to vaccines / vaccine ingredients including PEG/ Polysorbate